
New Clinical Fibromyalgia Diagnostic Criteria – Part 1.

To answer the following questions, patients should take into consideration

- how you felt the **past week**,
- while taking your current therapies and treatments, and
- exclude your pain or symptoms from other known illnesses such as arthritis, Lupus, Sjogren's, etc.

Determining Your Widespread Pain Index (WPI)

The WPI Index score from Part 1 is between 0 and 19.



Count up the number of areas checked and enter your Widespread Pain Index or WPI score score here

Symptom Severity Score (SS score) - Part 2a.

Indicate your level of symptom severity over the past week using the following scale.

Fatigue		Waking unrefreshed		Cognitive symptoms		
	0 = No problem1 = Slight or mild problems;generally mild or intermittent		0 = No problem1 = Slight or mild problems;generally mild or intermittent		0 = No problem1 = Slight or mild problems;generally mild or intermittent	
	2 = Moderate; considerable problems; often present and/or at a moderate level		2 = Moderate; considerable problems; often present and/or at a moderate level		2 = Moderate; considerable problems; often present and/or at a moderate level	
	3 = Severe: pervasive, continuous, life disturbing problems		3 = Severe: pervasive, continuous, life disturbing problems		3 = Severe: pervasive, continuous, life disturbing problems	

Tally your score for Part 2a (not the number of checkmarks) and enter it here

Symptom Severity Score (SS score)- Part 2b

Check each of the following OTHER SYMPTOMS that you have experienced over the past week?

	Muscle	pain		Nervousness		Loss/change in taste
	Irritable	bowel syndrome		Chest pain		Seizures
	Fatigue	/tiredness		Blurred vision		Dry eyes
	Thinkin	g or remembering problem		Fever		Shortness of breath
	Muscle	Weakness		Diarrhea		Loss of appetite
	Headac	he		Dry mouth		Rash
	Pain/cra	amps in abdomen		Itching		Sun sensitivity
□ Numbness/tingling □				Wheezing		Hearing difficulties
□ Dizziness □				Raynauld's		Easy bruising
☐ Insomnia ☐				Hives/welts		Hair loss
□ Depression □			Ringing in ears		Frequent urination	
□ Constipation □				Vomiting		Painful urination
☐ Pain in upper abdomen ☐			Heartburn		Bladder spasms	
□ Nausea □				Oral ulcers		
	up the mu u tallied:	umber of symptoms checked abor	ve.	Enter your score for Pa	art 2	b here
0 symp 1 to 10 11 to 2 25 or r	Give yourself a score of 1 Give yourself a score of 2		Now add Part 2a AND 2b scores, and enter This is your Symptom Severity Score (SS score), which can range from 0 to 12.			

What Your Scores Mean

A patient meets the diagnostic criteria for fibromyalgia if the following 3 conditions are met:

1a. The WPI score (Part 1) is greater than or equal to 7 <u>AND</u> the SS score (Part 2a & b) is greater than or equal to 5

OR

- **1b.** The WPI score (Part 1) is from 3 to 6 <u>AND</u> the SS score (Part 2a & b) is greater than or equal to 9.
- **2.** Symptoms have been present at a similar level for at least 3 months.
- **3.** You do not have a disorder that would otherwise explain the pain.

For example:

If your WPI (Part 1) was 9 and your SS score (Parts 2a & b) was 6, then you **would meet** the new FM diagnostic criteria.

If your WPI (Part 1) was 5 and your SS score (Parts 2a & b) was 7, then you **would NOT** meet the new FM diagnostic criteria.

*The new FM diagnostic criteria did not specify the number of "Other Symptoms" required to score the point rankings from 0 to 3. Therefore, we estimated the number of symptoms needed to meet the authors' descriptive categories of:

- 0 = No symptoms
- 1 = Few symptoms
- 2 = A moderate number
- 3 = A great deal of symptoms
- * Wolfe F, et al. Arthritis Care Res 62(5):600-610, 2010.

For information about Fibromyalgia Network, call our office Monday through Friday, 9:00 a.m. to 5:00 p.m. (PST) at (800) 853-2929 or visit us online at www.fmnetnews.com.

This survey is not meant to substitute for a diagnosis by a medical professional. Patients should not diagnose themselves. Patients should always consult their medical professional for advice and treatment. This survey is intended to give you insight into research on the diagnostic criteria and measurement of symptom severity for fibromyalgia.